

COMPOSITION :

Each 1 gm contains :

Amoxicillin (as amoxicillin trihydrate).....500 mg

**AMOXY 500 - 1KG****ANTIBIOTIC SOLUBLE POWDER****FOR ORAL ADMINISTRATION****INDICATIONS:**

It is freely water – soluble, suitable for all types of potable water.

Amoxicillin is a hemi-synthetic and a broad-spectrum penicillin, acting against bacteria GRAM-positive and GRAM-negative.

Levels of increasing resistance have been observed concerning E.coli, Salmonella typhimurium and Pasteurella haemolytica.

Calves: Respiratory infections caused by Pasteurella spp. and Salmonella dublin.

Gastrointestinal infections caused by E.coli. Infections caused by Salmonella dublin.

Poultry: Respiratory infections caused by E.coli, Haemophilus spp, Pasteurella spp and Salmonella spp. gastrointestinal infections caused by Salmonella spp.

CONTRAINDICATIONS:

Do not administer to animals with hypersensitivity to penicillins and cephalosporins, to animals with serious renal disorders and other animal species not listed at the indications. It is recommended to use gloves and mask during the administration of the product.


INTERACTION WITH OTHER MEDICINAL PRODUCT:

Do not administer simultaneously with effective bacteriostatic antibiotics such as tetracyclines, etc

WITHDRAWAL PERIOD:

Calves: Meat: 2 days, Milk: 2 days.

Poultry: Meat: 1 day, Egg: 0 day.

**DOSAGE AND ADMINISTRATION:**

Calves: 75 g per 100 liters of milk substitute or potable water twice a day, for 5 consecutive days (10 mg amoxicillin per kg of body weight /12 hours)

Poultry: The recommended dosage is 15 mg amoxicillin trihydrate per kg.

The total period of treatment should be for 3 consecutive days or in severe cases for 5 consecutive days. The medicated water must be replaced every 12 hours. The medicated milk replacer must be administered within the next 6 hours.

Keep it In original pack, Store in a dry, dark place below 25°C.

USED FOR

Calves & Poultry

