

COMPOSITION :

Each 1 gm contains :
Amoxicillin (as amoxicillin trihydrate).....200 mg



AMOXY 200 - 1KG

**ANTIBIOTIC SOLUBLE POWDER
FOR ORAL ADMINISTRATION**

INDICATIONS:

It is freely water – soluble, suitable for all types of potable water.
Amoxicillin is a hemi-synthetic and a broad-spectrum penicillin, acting against bacterias GRAM-positive and GRAM-negative. Levels of increasing resistance have been observed concerning E.coli, Salmonella typhimurium and Pasteurella haemolytica.
Calves: Respiratory infections caused by Pasteurella spp. and Salmonella dublin.
Gastrointestinal infections caused by by E.coli.
Infections caused by Salmonella dublin.
Poultry: Respiratory infections caused by E.coli, Haemophilus spp, Pasteurella spp and Salmonella spp. gastrointestinal infections caused by Salmonella spp.

CONTRAINDICATIONS:


Do not administer to animals with hypersensitivity to penicillins and cephalosporins, to animals with serious renal disorders and other animal species not listed at the indications.

INTERACTION WITH OTHER MEDICINAL PRODUCT:

Do not administer simultaneously with effective bacteriostatic antibiotics such as tetracyclines, etc

WITHDRAWAL PERIOD:

Calves: Meat: 2 days, Milk: 2 days.
Poultry: Meat: 1 day, Egg: 0 day.



DOSAGE AND ADMINISTRATION:

Calves: 75 g per 40 liters of milk substitute or potable water twice a day, for 5 consecutive days (10 mg amoxicillin per kg of body weight /12 hours)
Poultry: 150 - 300 g per 60 liters of potable water, for 6 consecutive days

The medicated water must be replaced every 12 hours. The medicated milk replacer must be administered within the next 6 hours.
It is recommended to use gloves and mask during the administration of the product, especially in case of known hypersensitivity to penicillin.

Keep it In original pack, Store in a dry, dark place below 25°C.

USED FOR

Calves & Poultry

