

treatment and prevention of coccidiosis in calves, goats, sheep, chickens (broilers and breeders) and other fowl such as turkeys, with activity against Eimeria spp., especially Eimeria tenella and Eimeria necatrix. It is a thiamine (vitamin B1) analogue and its pharmacological effect relies on competitive inhibition of thiamine uptake. Amprolium competitively inhibits the active transport of thiamine in isolated second generation schizonts of Eimeria spp., and in host intestinal cells. Amprolium indicated for coccidiosis caused by coccidia susceptible to amprolium (Eimeria spp.), or gastrointestinal infections for which it is therapeutically or prophylactically indicated to administer amprolium, in calves, goats, sheep and poultry.

CONTRAINDICATIONS:

Don't administer to animals with impaired liver and/or renal functions. Do not administer to turkeys before the age of 8 to 10 weeks.

WITHDRAWAL PERIOD:

Meat:Calves, goats, sheep: 3 days. Meat: Poultry & turkeys: 3 days.

Egg: Do not administer animals whose eggs are

intended for human consumption.

DOSAGE AND ADMINISTRATION:

CARLER S ALALAMA

-Be incorporated into the drinking water for 5 -7 days.

Poultry & turkeys: 1 Liter of AMPRO 250 in 5,000L of drinking water for 5-7 days.

Calves, sheep and goats: 1 mL of AMPRO 250 per 25 kg body weight through drinking water or milk for 5 days.

Always ask the veterinarian for the final dosage for your animal.

KEEP OUT OF REACH OF CHILDREN STORE IN CLOSE CONTAINER BELOW 25 °C

USED FOR



🎺 Poultry, calves, sheep & goat