

COMPOSITION:

Each 1000 gm contains : Acetylsalicylic acid500 Gm

Ascorbic Acid (Vitamin C)250 Gm



WATER SOLUBLE POWDER

INDICATIONS:

FOR ORAL ADMINISTRATION

Poultry: As an anti-stress factor, in heat strokes, post vaccinal reaction, in viral and acute bacterial diseases. Large Animal (Sheep, Goat, Camel): As a post operative pain killer.

In case of rheumatic pains, tendenitis, myositis and neuritis. Supportive treatment for reduction of pyrexia in combination with, appropriate anti-infective therapy, if necessary.

DESCRIPTION:

Acetylsalicylic acid is well-known, non-steroidal anti-inflammatory and one of the weak analgesics. Acetylsalicylic acid has an anti-inflammatory effect, a peripheral analgesic and antipyretic effect and it inhibits thrombocyte aggregation. The effectiveness of acetylsalicylic acid is due primarily to the inhibition of the enzyme cyclooxygenase, responsible for the formation of prostaglandins and thromboxane from arachidonic acid.

Prostaglandins are significant causes of inflammations, pain and fever.

Vitamin C (Ascorbic Acid) is involved in the conversion of folic acid to tetrahydrofolic acid and the conversion of proline to hydroxyproline which is essential to the formation of collagen.

CONTRAINDICATIONS:

Do not use in case of hypersensitivity to the active substance or in cases of gastrointestinal irritation and ulcers, chronic gastro-intestinal disorders, bronchospasm, liver function impairment or nephropathies.

Do not use in pregnant or lactating animal. Do not use in reminant less than 4 weeks of age.



Camel, Calves, goats and sheep:

2gm Twice daily per 100 kg body weight for 3 - 5 days. Poultry:

50gm twice daily per 100 litre of drinking water for 3 - 5 days.

1KG

WITHDRAWAL PERIOD:

Meat: Poultry, Camel, Calves, goats and sheep: 1 day.

Milk: Camel, Calves, goats and sheep: 1 day.

Egg: Poultry: 1 day.

Keep it In original container in a dry, dark place below 25°C

USED FOR

Poultry, Camel, Calves, goats and sheep