

COMPOSITION :

Each 1 gm contains :

Spectinomycin.....444.7 mg

Lincomycin.....222 mg



LINCO-SPECT

WATER SOLUBLE POWDER**FOR ORAL ADMINISTRATION****INDICATIONS:**

Gastrointestinal and respiratory infections caused by lincomycin and spectinomycin sensitive micro-organisms, like *Campylobacter*, *E. coli*, *Mycoplasma*, *Salmonella*, *Staphylococcus*, *Streptococcus*, and *Treponema spp.*, in Calves, Lambs, Kids, poultry, and turkeys.

The combination of lincomycin and spectinomycin acts additive and in some cases synergistic. Spectinomycin acts bacteriostatic or bactericidal, depending on the dose, against mainly Gram -ve bacteria like *Campylobacter*, *E. coli*, *Mycoplasma* and *Salmonella spp.* Lincomycin acts bacteriostatic against mainly Gram +ve bacteria like *Mycoplasma*, *Staphylococcus*, *Streptococcus* and *Treponema spp.* Cross resistance of lincomycin with macrolides can occur.

SIDE EFFECTS:

In the event of overdose in pigs, a change in the consistency of the faeces (soft faeces and/or diarrhoea) may be observed. In chickens treated at several times the recommended dose, enlargement of the caecum and abnormal caecum content was observed.

**DOSAGE AND ADMINISTRATION:****For oral administration:**

Calves, Lambs and Kids : 1.50 gram per 100 kg body weight for 5 days.

Poultry : 150 gm per 200 drinking water for 5 - 7 days.

WITHDRAWAL PERIOD:

Calves, Lambs and Kids: Meat: 14 days.

Poultry: Meat: 5 days.

Egg: Do not administer for layers producing eggs for human consumption.

Keep it In the Close container below 25°C

USED FOR

Calves & Poultry

