



COMPOSITION :

Each 1 KG contains :

Lincomycin base 500 gm which is equivalent to 567gm Lincomycin hydrochloride.

LINCOMED 500

**ANTIBIOTIC SOLUBLE POWDER
FOR ORAL ADMINISTRATION**

INDICATIONS:

Treatment against necrotic enteritis caused by pathogen agent susceptible to lincomycin, in particular clostridium perfringens.

SPECIAL WARNINGS:

Presence of indicated pathogens has to be shown. Bad use of the product may cause increasing prevalence bacteria resistant to lincomycin and reduce efficacy of the treatments with other lincosamides, macrolides and streptogramins due to possible appearing of a cross resistance.

Special precaution to be taken by person administering the veterinary medicine product to animals:

Protective gloves are suggested. Do not eat, drink or smoke during manipulation. Well wash hands after use. Persons with known hypersensitivity to lincomycin have to avoid contact with veterinary medicine.

INTERACTION WITH OTHER VET. MEDICINES AND OTHER FORMS OF INTERACTION:

Caoline reduces the absorption of lincomycin, if it is administered together.

In case of need the administration of both them, it is necessary to divide treatments at least of 2 hrs. The use together lincosamides and macrolides antibiotics give a decreasing of efficacy of both due to the steric abstraction of molecule in the seat of attack of the other. The lincomycin decreases the bacteria activity of aminoglycosidici. Do not administer with penicillin and cephalosporin.



DOSAGE AND ADMINISTRATION:

50gm LINCOCIN per 200 liters of drinking water for 5 days (correspond to 10/20 mg of product per kg b.w). Do not mix with solid feed.

To ensure a correct dosage, body weight should be determined as accurately as possible.

The intake of medicated water depends on the clinical condition of animals.

The medicated water should be the only source of drinking water throughout the treatment period. Medicated water should be refreshed every 24 hrs.

WITHDRAWAL PERIOD:

Meat and viscera : 5 days

**Keep it n cool and dry place.
Do not expose to direct heating sources.**

USED FOR

Poultry

