

**COMPOSITION :**

Each Bolus contains :

|                    |         |
|--------------------|---------|
| Sulphadiazine..... | 1000 mg |
| Trimethoprim.....  | 200 mg  |

**SULPHAPRIM-Bolus****SCOUR HALT ANTI-DIARRHETIC BOLUS**  
for Oral Administration**INDICATIONS:**

Sulphaprim Boluses are primarily indicated for oral use for the treatment of bacterial scours but may also be used for the treatment of acute salmonellosis and bacterial pneumonia.

Together the active ingredients produce a double blockage of folic acid synthesis in bacteria giving a level of activity many times greater than that obtained from either drug alone.

In vitro Sulphaprim Bolus is effective against most common Gram-positive and Gram-negative bacteria including

*Escherichia coli*, *Klebsiella* spp, *Pasteurella* spp, *Salmonella* spp, and *Staphylococcus* spp.,

The boluses are particularly effective against *Escherichia coli* and *Salmonella* spp infections.

**SIDE EFFECTS:**

As with all veterinary products, some unwanted effects can occur from the use of sulfadiazine and trimethoprim boluses.

Always consult a veterinary physician or animal care specialist for medical advice before use. Common side effects include Hypersensitivity reactions and hematologic effects, including anemias, thrombocytopenia, or leukopenias and diarrhea. For a comprehensive list of all possible effects, consult a veterinary physician. If any symptom persists or gets worse, or you notice any other symptom, then please seek veterinary medical treatment immediately.

**DOSAGE AND ADMINISTRATION:****Calves, Sheep and Goat:**

One bolus per 40 kg bodyweight daily.

This gives 30 mg of combined active ingredients per kg. Treatment should be repeated daily until two days after the symptoms have resolved, but in cases of salmonellosis and bacterial pneumonia, treatment should be continued for 5 consecutive days.

Treatment must not be continued for more than 5 days.

**WITHDRAWAL PERIOD:**

Sheep: Meat: 14 days.

Calves: Meat: 14 days.  
use for human consumption.

Goat: Meat: 14 days.  
Milk: Not intended for

**CONTRA-INDICATIONS:**

Administration in case of renal or hepatic insufficiency, chronic infections, abscess formation.

**Keep it in the original packing  
in a dry, dark place below 25°C**

**USED FOR**

Calves, Sheep and Goat.

