

COMPOSITION :

Each 1 ml contains :
 Toltrazuril.....25 mg



TOLTRACOX - 1L-250ml

ANTICOCCIDIAL SUSPENSION

FOR ORAL ADMINISTRATION

INDICATIONS:

For the treatment of coccidiosis caused by:
 Chicken (pullets and breeders): *Eimeria acervulina*,
E. brunetti, *E. maxima*, *E. necatrix* and *E. tenella*.
 Turkeys: *Eimeria adenoides*, *E. meleagrimitis*.
 Calves: For the prevention of clinical signs of coccidiosis
 and reduction of coccidia shedding in housed calves
 replacing cows producing milk for human consumption
 (dairy cows) on farms with a confirmed history of
 coccidiosis caused by *Eimeria bovis* or *Eimeria zuernii*.
 Lambs: For the prevention of clinical signs of coccidiosis
 and reduction of coccidia shedding in lambs on farms
 with a confirmed history of coccidiosis caused by
Eimeria crandallis and *Eimeria ovinoidalis*.

DESCRIPTION:

Toltrazuril is an anticoccidial of the triazinetrione group,
 active against *Eimeria* spp. Its activity affects the
 intracellular development stages of the parasites without
 affecting the extracellular stages of the parasites.
 At parasite level, toltrazuril decreases the enzymatic
 activity of the respiratory chain, causing inflammation
 of the endoplasmic reticulum and Golgi apparatus,
 perinuclear space modifications and alteration of
 division of the nucleus.

In chickens and turkeys, toltrazuril is absorbed at rate
 of at least 50%. Distribution is higher in liver and kidney.
 The active substance is rapidly metabolised and the
 main metabolite is characterised as a toltrazuril sulfone.

WITHDRAWAL PERIOD:

Poultry : Meat : 16 Days. Turkeys: Meat : 18 Days.
 Calves: Meat : 63 Days. Lambs: Meat : 42 Days.
 Eggs: Not used for Layers producing eggs for human
 consumption.



DOSAGE AND ADMINISTRATION:

- Poultry & Turkeys:

The recommended dose rate is 7 mg toltrazuril per kg
 of body weight (equivalent to 28 mL of medicinal
 product per 100 kg of body weight) daily given for
 2 consecutive days.

- Calves:

Each animal should be treated with a single oral dose
 of 15 mg toltrazuril/kg body weight corresponding to
 6.0 mL oral suspension per 10 kg body weight.

- Lambs:

Each animal should be treated with a single oral dose
 of 20 mg toltrazuril/kg body weight corresponding to
 0.8 mL oral suspension per kg body weight

**Keep it In original
 container below 25°C.**

USED FOR

Poultry, Calves, Lambs

