

COMPOSITION :

Each 1 gm contains :

Doxycycline Hyclate.....100 mg

Tylosin Tartrate.....100 mg

TYLODOX 100-100**WATER SOLUBLE POWDER****FOR ORAL ADMINISTRATION****INDICATIONS:**

It is freely water – soluble, suitable for all types of potable water.

It is used to indicate Gastrointestinal and respiratory infections caused by tylosin and doxycycline sensitive micro-organisms, like: Bordetella , Campylobacter, Chlamydia , E.coil , Haemophilus, Mcoplasma , Staphylococcus, Treponema spp , Rickettsia, Salmonella, Streptococcus and Pasteurella . , in cattle, goats , poultry and sheep

WITHDRAWAL PERIOD:

Cattle, Goats and sheep: Meat: 14 Days.

Milk: Do not use in animals producing milk for Human Consumption.

Poultry: Meat: 8 Days.

Egg: Not to use in Layers producing eggs for Human Consumption

CONTRA-INDICATIONS:

Do not administer to animals with hypersensitivity to tylosine or doxycycline, to animals with serious renal disorders and other animal species not listed at the indications

**DOSAGE AND ADMINISTRATION:****For oral administration:**

Cattle: Twice daily 10 gram per 200 kg body weight for 3-5 days.

Goats and sheep: Twice daily 10 gram per 200 kg body weight for 3-5 days.

Poultry: 100 gram per 100-200 litre drinking water for 3-5 days.

The medicated water must be replaced every 12 hours.

The medicated milk replacer must be administered within the next 6 hours

Keep it In the Close container below 25°C

USED FOR

Poultry, Cattle, Sheep and Goat 