



TYLODOX 200-200

WATER SOLUBLE POWDER FOR ORAL ADMINISTRATION

INDICATIONS:

It is freely water – soluble, suitable for all types of potable water. It is used to indicate Gastrointestinal and respiratory infections caused by tylosin and doxycycline sensitive micro-organisms, like Bordetella , Campylobacter , Chlamydia , E.coil , Haemophilus, Mcoplasma , Pasteurella ,Staphylococcus, Streptococcus , Rickettsia, Salmonella, and Treponema spp. in Calves, Kids, Lambs and Poultry.

WITHDRAWAL PERIOD:


Calves: Meat: 14 Days,
Lambs: Meat: 14 Days.
kids: Meat: 14 Days.
Poultry: Meat: 8 Days,
Egg: Do not use in layers producing eggs for human consumption.

CONTRA-INDICATIONS:

Do not administer to animals with hypersensitivity to tylosine or doxycycline, to animals with serious renal disorders and other animal species not listed at the indications

COMPOSITION :

Each 1 gm contains :
Doxycycline Hyclate.....200 mg
Tylosin Tartrate.....200 mg



DOSAGE AND ADMINISTRATION:

For oral administration:

Calves: Twice daily 5 gram per 200 kg body weight for 3-5 days.

Lambs and Kids: Twice daily 5 gram per 200 kg body weight for 3-5 days.

Poultry: 50 gram per 100-200 litre drinking water for 3-5 days.

The medicated water must be replaced every 12 hours.

The medicated milk replacer must be administered within the next 6 hours.

**Keep it In the original packing,
Store in a dry, dark place below 25°C**

USED FOR

Poultry, Calves, Lambs and Kids.

