

**COMPOSITION :**

Each 1 Liter contains :

Vitamin A.....	50,000,000 I.U.
Vitamin D3.....	10,000,000 I.U.
Vitamin E.....	10,000 I.U.
Vitamin C.....	50,000 Mg.
Vitamin K.....	1.250 Mg.

**VITAVET AD3ECK - 1 L****VITAMIN (A+D3+E+C+K) LIQUID  
FOR ORAL ADMINISTRATION****INDICATIONS:**

The product is recommended as a growth stimulant in the young stock of horses, cattle, sheep, goats, swine, poultry (hen, turkey, guinea-fowl, quail, pheasant, peacock, goose, duck, pigeon) dogs and cats, as well as for the improvement of the proliferation in breeder flocks. Vitamin AD3ECK can be also used as antistress treatment; if associates with antibiotics, it can be used in the prevention of some infectious or parasitic diseases (salmonellosis, colibacillosis, coccidiosis etc.). Vitamin AD3ECK can also be used as support for the convalescent animals or in case of disturbances of the mineral, fatty acid and carbohydrate metabolism.

**CONTRA INDECACTION:**


Do not use with other product containing vitamin A,D3, E,C, K.

**SIDE-EFFECTS:**

No side effects observe if not exceeding the recommended dose. Overcoming the marked doses or prolonged administration may produce adverse effects or poisoning by aggregation manifested by anorexia, color of skin or mucous membranes, anemia, leukopenia, splenic and hepatic hypertrophy.

**WITHDRAWAL PERIOD:**

Egg: 0 Days. Meat: 0 Days. Milk: 0 Days.

**DOSAGE AND ADMINISTRATION:**

- Horses , Camels and Cattle:  
1mL per 80KG bodyweight into the drinking water for 3 - 5 days,
  - Calves and adult Sheep, Goats :  
1mL per 40KG bodyweight into the drinking water for 3 - 5 days.
  - Poultry (hens, turkeys, guinea-fowls, quails, pheasants, peacocks, geese, ducks, pigeons):  
1 litre per 1,000 - 2000 litre drinking water for 3-5 days.
- \*\* The dose and repetition can be changed according to veterinary Doctor

**ADVICE ON CORRECT ADMINISTRATION:**

In animals refusing water, the product is administered using a syringe without needle, at mouth comisura.

**Keep it In a close container below 25°C**

**USED FOR**

Horses, Camels, Cattle, Sheep, Goats, Poultry.