

dystrophy, exudative diathesis, decreased hatchabillty in eggs, infertility problems) in calves, cattle, goats, poultry and sheep.

Prevention of iron intoxication after administration of iron.

DESCRIPTION:

Vitamin E is a fat-soluble intracellular antioxidant. involved in stabilising unsaturated fatty acids. The main antioxidant property is preventing formation of toxic free radicals and oxidation of the unsaturated fatty acids in the body. These free radicals can be formed in periods of disease or stress in the body. Selenium is an essential nutrient for animals. Selenium is a component of the enzyme glutathione peroxidase, which plays an important role in protection of cells by destroying oxidising agents like free radicals and oxidated unsaturated fatty acids

CONTRAINDICATIONS:

No undesirable effects are to be expected when the prescribed dosage regimen is followed.

WITHDRAWAL PERIOD:

Poultry & Turkeys: Meat: 0 Day Eggs: 0 Day. Calves, goats and sheep: Meat:0 Day. Cattle & Camels: Meat: 0 Day Milk: O Day.

DOSAGE AND ADMINISTRATION:

For Oral Use.

Poultry & Turkeys:

1 liter per 4 000 litre drinking water for 3-5 days. Chicks (< 21 days):

1 Liter per 1 000 litre drinking water for 5- 10 days. Calves, goats and sheep:

10 mL per 50 kg. body weight for 3- 5 days.

PACK SIZE: 1LITER

Cattle & Camels:

10 mL per 80 kg. body weight for 3 to 5 days. Treatment may be repeated if considered necessary. Mixed water should be used in 24 hour.

> Keep it In original packing in a dry, dark place below 25°C

USED FOR

Poultry, Cattle, Calves, goats and sheep