



VITAVET FERTIL AID

VITAMINS, MINERALS & AMINO ACIDS POWDER

for Oral Administration

COMPOSITION :

Each 5 KG contains :

Vitamin A.....	285 M.I.U
Vitamin D3.....	200 M.I.U
Vitamin E.....	125 gm
Sodium Selenite.....	0.012 gm
Vitamin K3.....	0.25 gm
Vitamin B1.....	2.5gm
Vitamin B2.....	2.5 gm
Vitamin B6.....	2.5 gm
Ferrous Gluconate.....	0.1440 gm
L-Valin.....	2.5 gm
Copper Sulphate.....	0.04 gm
Cobalt Sulphate.....	1.25 gm
Alanine.....	2.5 gm
L-Arginine.....	1.25 gm
L-Aspartic Acid.....	4.25 gm
L-Cysteine.....	1.1 gm
L-Glutamic acid.....	0.008 gm
L-Histidine.....	0.75 gm
L-Proline.....	1.375 gm
L-Serine.....	2.125 gm
L-Threonine.....	2 gm
L-Tryptophan.....	0.7gm

Vitamin B12.....	0.005 gm
Niacin.....	12.5 gm
Vitamin C.....	50 gm
Folic Acid.....	12.5gm
Manganese Sulphate.....	100 gm
Magnesium Sulphate.....	314.5 gm
Zinc Sulphate.....	434.75 gm
Ferric Ammonium Citrate.....	0.1 gm
Potassium Iodide.....	0.4 gm
Co Q10.....	5 gm
D-Biotin.....	0.75 gm
Calcium Phosphate.....	69.35 gm
Calcium Gluconate.....	151.35 gm
Calcium Acetate.....	65.75 gm
L-Leucine.....	5 gm
L-Isoleucin.....	2.5 gm
DL-Methionine.....	1 gm
L-Phenylalanine.....	1.62 gm
L-Tyrosine.....	0.35 gm
L-Glycine.....	10 gm
L-lysine.....	3.75 gm
L-carnitine.....	1 gm
Ferrous Amino Acid Chalet.....	0.162 gm



DOSAGE AND ADMINISTRATION:

Be incorporated into the drinking water for 5-7 days, the following doses per animal per day:

Horses, Camels, Cattle, Sheep and Goat:

1 gram per 2 - 4 litre drinking water for 5 - 7 days.

Poultry :

2 gram per 5 litre drinking water for 5 - 7 days.

WITHDRAWAL PERIOD:

Meat: 0 days.

Milk: 0 days.

Egg: 0 days.

Keep it in a dry, dark place below 25°C

USED FOR



Horses, Camels, Cattle, Sheep and Goat.



INDICATIONS:

used to maintain normal vital process, development and production.

Also Increment of reproductive performance in both males and females of different animals. Improves quality, fertility and hatchability rates, and increases the eggproduction. In hot days, it combats anorexia and increases feed and water intake.

After treatment of coccidiosis, worm infestation bacteria also Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.