

**COMPOSITION :**

Each 1 GM contains:

Vitamin K3.....125 MG



# VITAVET K3 125

## VITAMIN K3 POWDER for Oral Administration

### INDICATIONS:

In Poultry: Prevention & treatment of Vitamin K3 deficiency, Haemorrhagic diseases (Gumboro, Coccidiosis, heavy parasitic infestation), Before debeaking and vaccination, Egg blood spots, Stress conditions, Reduce mortality rate in the embryonic stages.

In Horse, Cattle, Camel, Sheep and Goat : Epistaxis in racing horses, Jaundice of newborn, Anaemia, haemorrhages & heavy metal intoxication. Hepatic and intestinal disorder, Preventive bleeding measure in delivery & surgery.

### DESCRIPTION::

Used to repair the deficiency of Vitamin K3 causes a reduction in the prothrombine content of the blood which prevents blood clotting in haemorrhage cases.

### NOTE:

After diluting the product into the potable water the final solution remains stable for 24 hours at temperature (15-25C).

After diluting the product into the milk, the solution should be immediately consumed.

### WITHDRAWAL PERIOD:

Meat: Poultry & turkeys : 0 days.

Egg: Poultry & turkeys : 0 days.



## DOSAGE AND ADMINISTRATION:

### By oral administration.

Be incorporated into the drinking water for 3-5 days, the following doses per animal per day:

**Adult horses , camels, cattle, Sheep and goat:** 8-10 gm of vitamin K3 per 150 kg body weight added to feed or in drinking water for 5-8 days.

**Poultry (hens, turkeys, guinea-fowls, quails, pheasants, peacocks, geese, ducks, pigeons):** 4-5 gm of VITAVET K3 per 10 liter of drinking water for 3 - 5 days.

**Double the doses in severe cases. Or as directed by the registered veterinary physician.**

**Consumption may vary depending on factors like age, state of health, breed, husbandry system.**

**Keep it in the original packing in  
a dry, dark place below 25°C**

## USED FOR

Horses, cattle, sheep, goats, poultry