

COMPOSITION :

Each 1 ml contains :

Imidocarb Dipropionate120mg

**IMOCARB - 10ml-50ml
100ml**

Long duration of activity ANTI PROTOZOAL.

Injection Solution**INDICATIONS:**

Used to control **Babesiosis** in cattle and dogs.
As well as treatment and prevention of
Anaplasmosis in cattle and **Ehrlichia canis**
infections in dogs.

CONTRA-INDICATIONS:

Do not administer **intravenously** in cattle.
Do not use in cases of **hypersensitivity** to the
active substance or to any of the excipients.

DESCRIPTION:

Imidocarb is an antiprotozoal derived from
carbanilide,
Which act directly on the glycolysis of the parasite
and as an inhibitor of topoisomerase II, blocking
DNA replication.

Its spectrum of action includes:**- Cattle:**

Babesia argentina, *B. bigemina*, *B. bovis*,
B. divergens, *Anaplasma marginale*.

- Dogs:

Babesia canis, *B. gibsoni*, *B. vogelli*.

WITHDRAWAL PERIOD:

Slaughter of cattle for **meat** allowed no earlier
than **21 days**,
for **milk** consumption no earlier than **6 days**.

**DOSAGE AND ADMINISTRATION:****(For Subcutaneous injection Only).****Cattle:**

For Babesiosis: 1ml/100 Kg Body Weight.

For Anaplasmosis or mixed infections anaplasma and
babesia: 2.5 ml/100 kg.

Dogs:

For Babesiosis: 0.25 ml/10 Kg Body Weight.

For Ehrlichiosis or mixed infections due to *Ehrlichia*
and *Babesia*: 0.5 ml/ 10 kg.

2 doses at an interval of 14 days.

**Keep it in the original packing at temperatures
not exceeding 25°C, protected from light.**

USED FOR

Cattle, Dogs