

COMPOSITION :

Each 1 ml contains :

Ivermectin.....10mg

**IVECTINE - (50ml , 100ml)**

ANTI-PARASITE MEDICATION FOR PARASITIC DISEASES.

Injection Solution**INDICATIONS:**

Treatment and control of the following harmful parasites of CATTLE: Gastrointestinal Roundworms, *Ostertagia lyrata*, *Ostertagia ostertagi*, *Cooperia oncophora*, *Cooperia pectinata*, *Cooperia punctata*, *Haemonchus placei*, *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Bunostomum phlebotomum*, *Oesophagostomum radiatum*, *Strongyloides papillosus*, *Nematodirus helvetianus*, *Nematodirus spathiger*, *Trichuris* spp., Lungworms, *Dictyocaulus viviparus*, Eye Worms, *Thelazia* spp , Warbles, *Hypoderma bovis*, *H. lineatum*, Mange Mites, *Psoroptes ovis*, *Sarcoptes scabiei* var. *bovis*, Sucking Lice *Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*. Also be used as an aid in the control of biting lice (*Damalinea bovis*) Mange mite *Chorioptes bovis*, but complete elimination may not occur.

SHEEP: Gastrointestinal Roundworms, *Ostertagia circumcincta*, *O. trifurcata*, *Haemonchus contortus*, *Trichostrongylus axei*, *T. colubriformis*, *T. vitrinus* *Cooperia curticei*, *Oesophagostomum columbianum*, *O. venulosum*, *Nematodirus filicollis*, *Chabertia ovina*, *Trichuris ovis*, Lungworms, *Dictyocaulus filaria* *Protostrongylus rufescens*, Nasal Bots, *Oestrus ovis*, Mange Mites, *Psoroptes ovis*.

WITHDRAWAL PERIOD:

Cattle (meat & Offal) : 49 days.

Sheep (meat & offal) : 37 days.

Sheep or Cattle (milk) : Do not use in sheep or cattle producing milk for human consumption.

**DOSAGE AND ADMINISTRATION:****(For Subcutaneous injection Only).**

for Cattle and Sheep given at the recommended dosage of **0.2 mg per kg** it controls re-infection with: Barbers pole worm - *Haemonchus placei* 14 Days. Small intestinal worm - *Cooperia* spp. 14 Days. Hairworm - *Trichostrongylus axei* 14 Days. Brown stomach worm - *Ostertagia ostertagi* 21 Days. Nodular worm - *Oesophagostomum radiatum* 21 Days. Lungworm - *Dictyocaulus viviparus* 28 Days. **The timing of treatment should be based on epidemiological factors and should be customised for each individual farm.**

CONTRA-INDICATIONS:

Do not use intramuscularly or intravenously for Cattle and Sheep. it is has been formulated specifically for use in these target species. Do not use in other species as severe adverse reactions.

Keep it In original pack, Store in a dry, dark place below 25°C.

USED FOR

Cattle, sheep

