

COMPOSITION :

Each 1 Liter contains :

BETAINE HCL.....	20,000 Mg
L-CARNITINE.....	500 Mg
DL-METHIONIN.....	10,000 Mg
VITAMIN B12.....	10,000 µg
NIACIN.....	10,000 Mg
INOSITOL.....	10,000 Mg
D-BIOTIN.....	500 Mg
SORBITOL.....	10,000 Mg
FOLIC ACID.....	1,000 Mg
Di-Methyl Glyceride.....	50,000 Mg

VITAVET LIVER AID**VITAMINS & AMINO ACIDS LIQUID**
for Oral Administration**INDICATIONS:**

LIVER AID guards the liver against various toxins, chemical drugs and toxic effect of the feed contaminants.

LIVER AID also helps in rapid regeneration of liver tissues and cells.

LIVER AID stimulates sluggish liver parenchyma cells and restores liver functions. LIVER AID maintains proper feed intake, digestion, absorption of nutrients.

DESCRIPTION:

LIVER AID contains Vitamins and Amino Acids where they are working together to prevent and treat livers. Betaine prevents and treats fatty liver in a moderate high-dietary-fat model in Animals.

L-carnitine supplementation improves liver function by reducing histological steatosis.

Methionine is used to prevent liver damage in acetaminophen (Tylenol) poisoning. It is also used for increasing the acidity of urine, treating liver disorders, and improving wound healing.

Niacin inhibits fatty acid flux from adipose tissue to liver, reduces hepatic triglyceride synthesis and increases hepatic lipid oxidation. Thus, niacin may have a role in the regulation of liver fat content.

Folic acid (folate) is a type of B vitamin that is key for cell growth and can work for liver regeneration.

**DOSAGE AND ADMINISTRATION:**

Add 1mL per 3 liter of drinking water.

Treatment maybe repeated if considered necessary.

The dose and repetition can be changed according to veterinary Doctor.

WITHDRAWAL PERIOD:

Milk : 0 Days. Meat : 0 Days. Egg: 0 Days.

SIDE-EFFECTS:

No undesirable effects are to be expected when the prescribed dosage regimen is followed.

KEEP OUT OF REACH OF CHILDREN
STORE IN CLOSE CONTAINER BELOW 25 °C

USED FOR

Poultry, Camels