

COMPOSITION :

Each 1 ml contains :

Albendazole.....25 mg.

WORMVET 2.5%**DEWORMER SUSPENSION**
For Oral Administration**INDICATION:**

A broad spectrum multi-purpose anthelmintic for the control of mature and developing immature forms of gastrointestinal roundworms, lungworms, tapeworms and adult liver fluke in cattle and sheep.

The product is also ovicidal against fluke and roundworm eggs.

In cattle it is active against the following species:

(Round worms: *Ostertagia*, *Haemonchus*, *Trichostrongylus*, *Bunostomum*, *Nematodirus*, *Oesophagostomum*, *Cooperia* and *Strongyloides* spp). It is usually effective against inhibited larvae of *Cooperia* and *Ostertagia*, Lungworms: *Dictyocaulus viviparus*, Tapeworms: *Moniezia* spp., Adult liver fluke: *Fasciola hepatica*.

In sheep it is active against benzimidazole susceptible strains of the following species: Roundworms: *Teladorsagia* (*Ostertagia*), *Haemonchus*, *Trichostrongylus*, *Nematodirus* (including *N. battus*), *Chabertia* and *Oesophagostomum*. It is usually effective against inhibited larvae of *Ostertagia*, Lungworms: *Dictyocaulus filaria*, Tapeworms: *Moniezia* spp., Adult liver fluke: *Fasciola hepatica*. The product is ovicidal and will kill fluke and roundworm eggs, thus reducing pasture contamination.

SIDE-EFFECT:
No side effects observe if not exceeding the recommended dose.

WITHDRAWAL PERIOD:

Cattle: meat: 15 days, milk: 5 Days
Sheep and Goat: meat: 15 days, milk: 5 days.

Do not use in Goat producing milk for human consumption.

**DOSAGE AND ADMINISTRATION:**

To ensure administration of a correct dose, body weight should be determined as accurately as possible; accuracy of the dosing device should be checked. Do not mix with other products.

Cattle:

(Worm dose): For the control of roundworms, lungworms, tapeworms and fluke and roundworm eggs.

Dosage: 75mg albendazole /kg.

(Fluke and worm dose): For the additional treatment of adult liver fluke (chronic fascioliasis) in cattle.

Dosage: 10 mg albendazole/ kg.

Sheep and goat:

(Worm dose): For the control of roundworms, lungworms, tapeworms, fluke and roundworm eggs.

Dosage: 5 mg albendazole / kg.

(Fluke and Worm Dose): For the additional treatment of adult liver fluke (chronic fascioliasis) in sheep.

Dosage: 75 mg albendazole / kg.

**Keep it in the original packing
in a dry, dark place below 25°C**

USED FOR

Cattle, Sheep and Goat.

