

thereby preventing infections in the affected area. PVI 10% is for external use only, the affected area should be clean and dry before application , you must wash your hands thoroughly before and after applying this disinfectant .

this medicine should be used regularly to get the most benefit from it . do not use more than you need as it will not clear condition faster and some side effects may be increased you can help this medication wark better by keeping the affected areas clean. this medicine may cause side effects like skin irritation, redness, or itching for animals at the site of application. Please consult your veterinarians if these side effects persist for a longer duration. Avoid direct contact with eyes.In case of direct contact, wash eyes with water. You should stop giving to animal straight away if any signs of an allergic reaction. Signs of this include rash, sewlling of the lips, threat, or face. swallowing or breathing problems, dizzy or faint and nausea.

BINIFETS OF PVI 10%:

In treatmentand prevention of wound infection PVI 10% is a versatile antiseptic that is used for the treatment and prevention of infection. PVI 10% kills and prevents the growth of infection causing microbes. therapy preventing abrasions, cuts, and wounds or any break in the skin from getting infected. the antiseptic effect is caused due to the slow release of iodine. keep the affected area clean and ues the medicine as prescribed by your doctor.

HOW TO USE PVI 10%:

FOR EXTERNAL USE:

Use it as advised by veterinarian or check the lable for directions before use.

HOW PVI 10% WORK:

PVI 10% is anantiseptic applied on skin, which is infected or is likely to get infected. it works by slowly releasing iodine which kills or prevents the growth of innfectious microorganisms.

* If you miss a dose of PVI 10%, skip it and continue with your normal schedule. Do not double the dose.

SIDE EFFECTS:

Most side effects do not require any medical attention and dissppear as your body adjusts to the medicine. Consult your veterinarian doctor. Application site reactions (burning, irritation, itching and redness)

KEEP OUT OF REACH OF CHILDREN
STORE IN A DRY, DARK PLACE BELOW 25 °C

USED FOR

Animals