

# EPROCTINE

## EPRINOMECTIN INJECTION SOLUTION

### **COMPOSITION :**

Each 1 ml contains:

Eprinomectin ..... 20 mg  
Excipients q.s ..... 1 ml

### **INDICATIONS :**

For the treatment and control of internal and external parasites susceptible to eprinomectin, including gastrointestinal and pulmonary nematodes, sucking lice, mange mites, and other ectoparasites as specified in the package leaflet.

### **WITHDRAWAL PERIOD:**

Cattle: Meat and offal: 63 days.

Milk: zero hours.

Sheep: Meat and offal: 42 days.

Milk: zero hours.

Goats: Meat and offal: 42 days.

Milk: zero hours.

*Pregnant women and women of childbearing age should avoid exposure to this product.*

**Once opened, use within  
6 months, Store below 30°C.**



## DOSAGE AND ADMINISTRATION:

### **(Subcutaneous injection only):**

#### **Dosage:**

Administer 0.2 mg eprinomectin per kg body weight, equivalent to 0.1 mL of product per 10 kg body weight.

For goats, the injected volume at any one site should not exceed 0.6 mL.

Determine the animal's body weight as precisely as possible to ensure correct dosing. Verify the accuracy of the dosing device before use.

## USED FOR

Cattle, sheep, and goats.