

COMPOSITION

Each 1 ml contains:

Vitamin B ₁ (hydrochloride)	0.10 mg
Vitamin B ₂ (as sodium phosphate)	0.04 mg
Vitamin B ₆ (hydrochloride)	0.10 mg
Vitamin B ₁₂	0.05 µg
Nicotinamide	1.50 mg
d-Panthenol	0.10 mg

ELECTROLYTES:

Calcium chloride hexahydrate	0.23 mg
Magnesium sulphate · 7 H ₂ O	0.29 mg
Potassium chloride	0.20 mg

AMINOACIDS AND NUTRITIONAL AGENTS:

Anhydrous dextrose	45.46 mg
L-Arginine hydrochloride	0.025 mg
L-Cysteine Hcl monohydrate	0.01 mg
Monosodium glutamate	0.04 mg
L-Histidine HCl monohydrate	0.01 mg
L-Isoleucine	0.01 mg

L-Leucine	0.04 mg
L-Lysine hydrochloride	0.03 mg
L-Methionine	0.01 mg
DL-Phenylalanine	0.03 mg
L-Threonine	0.02 mg
DL-Tryptophan	0.01 mg
DL-Valine	0.05 mg

EXCIPIENTS

Methyl parahydroxybenzoate	1.80 mg
Propyl parahydroxybenzoate	0.20 mg
Phenol	0.10 mg
Disodium edetate	0.15 mg
Sodium acetate · 3 H ₂ O	2.50 mg
Citric acid monohydrate (to adjust pH)	
Water for injection q.s. ad	1.0 ml

NUTRI-LYTE

INJECTION SOLUTION

INDICATIONS :

Intended as a supportive therapy in cases of dehydration, electrolyte imbalance, or hypoproteinaemia, and to maintain normal fluid and mineral balance during recovery or stress in animals.

CONTRAINDICATIONS AND WARNINGS:

Do not administer to animals with a known sensitivity to any component of the formulation. Use with caution in debilitated animals and ensure compatibility with concurrently administered solutions.

GENERAL PRECAUTIONS:

When administering intravenously, inject slowly and maintain strict aseptic technique. Use sterile equipment for each injection. Do not reuse partially used containers. Dispose of empty bottles and waste material safely in accordance with local regulations.

Store in the original, tightly closed container in a cool, dark location (between +5°C and +25°C). Do not freeze.



DOSAGE AND ADMINISTRATION:

Horses: Administer slowly by intravenous injection only.

Cattle : Administer slowly by intravenous, intraperitoneal, or subcutaneous injection.

Dogs and cats: Administer slowly by intravenous or subcutaneous injection.

Day-old chicks: Administer by subcutaneous injection.

Dosage:

Horses, and cattle: Up to 100 mL per 50 kg B.W

Foals, and calves: Up to 30 mL per 5 kg B.W

Dogs and cats: Up to 50 mL per 5 kg B.W

Day-old chicks: 0.5 - 1.0 mL per bird.

USED FOR

Horses, Cattle, Dogs, Cats& Chicks