



## COMPOSITION :

Each 100 ml contains:

Copper (Copper Gluconate).....	2,500	mg
Manganese (Manganese chloride) .....	4,800	mg
Zinc (Zinc oxide) .....	4,000	mg
Ortho phosphoric acid .....	332,250	mg
Calcium chloride .....	60,900	mg
Sodium chloride .....	19,100	mg
Magnesium oxide .....	16,500	mg
Crude Fiber 0.0%	Crude Ash 37.4%	
Crude Protein 0.0%	Crude Fat 0.0%	
Calcium 2.20%	Methionine 0.0%	
Phosphorus 10.50%	Lysine 0.0%	
Magnesium 1.00%	Sodium 0.75%	

# VITAVET M-PHO

## FEED ADDITIVES

**INDICATIONS :** *VITAVET M-PHOS* is a liquid complementary mineral feed supplying phosphorus, calcium, magnesium, and essential trace elements (copper, zinc, manganese) in a readily absorbable form.

Formulated for cattle, camels, sheep, goats, and poultry, it helps maintain mineral balance, fertility, growth, and performance under intensive or stressful conditions. It compensates for mineral losses during milk production, egg formation, growth, pregnancy, or heat stress, supporting:

- Calcium-phosphorus metabolism for strong bones and firm eggshells.
- Fertility, lactation, and feed utilization by maintaining digestive and rumen efficiency.
- Healthy immune function, resilience, and vitality during stress or high demand.

## RECOMMENDATION:

- Peak laying or breeding seasons, supports eggshell formation and reproductive performance.
- Feed or environmental stress – helps maintain mineral balance and feed intake under changing conditions.
- Post-antibiotic or vaccination recovery – aids mineral replenishment and recovery.
- High-temperature or transport stress – supports hydration and immune stability.

Note: Use under the supervision of a veterinarian or animal nutritionist for best results.



## DOSAGE AND ADMINISTRATION:

**(oral administration in drinking water):**

Cattle (adult): 75 – 100 mL/day for 3 – 5 days (repeat monthly or around calving).

Calves: 25 – 50 mL/day for 3–5 days.

Sheep/Goats (adult): 10 – 15 mL/day for 3 – 5 days.

Lambs/Kids: 5 – 10 mL/day for 3–5 days.

Poultry: 15 – 30 mL per 100 birds/day for 3 – 5 days, or 1-2L per 1000L of drinking water.

**WITHDRAWAL TIME:** Meat: 0 day, Egg: 0 day.

**SIDE EFFECTS:** No side effects when follow the dosage.

**store in original packing,  
Store in a dry, dark place below 30°C.**

## USED FOR

**Cattle, Camels, Sheep, Goats, Poultry**